

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.


'Inferences from Maxwell's theory concerning the motion of pure ether' (Wissenschaftliche Abhandl. B. III., p. 526, Wiedem. Am. Vol. LIII., p. 135-143).

M. I. PUPIN.

COLUMBIA UNIVERSITY, April 2, 1896.

A METHOD OF DETERMINING THE RELATIVE TRANSPARENCY OF SUBSTANCES TO THE RÖNTGEN RAYS.

The fact that the Röntgen rays have the power of dissipating the charge of a perfectly insulated electrified body was established by Professor J. J. Thompson,* and furnishes us one of the simplest methods of detecting the rays. This effect is the basis of a very simple method of making quantitative measurements of the intensity of the radiation. If we take a condenser and allow the Röntgen rays to fall upon it, we shall find that there is a very considerable diminution in its insulation resistance, and that the charge of the condenser is gradually dissipated. This is illustrated by the



curves A and B in the accompanying figure. A was obtained under the ordinary conditions. B was obtained when the Crookes tube was in action, and placed about six

inches from the wooden side of condenser. The curve A was determined before, and again immediately after the determination The two determinations of A were identical, showing that the effect of the Röntgen ray on the insulation disappeared with the cessation of the ray. In making these measurements a Nalder micro-farad condenser was used, the condenser being charged with a standard Clark cell. It is evident, therefore, that it is possible to compare the transparency of different substances by allowing the rays to pass through screens made of the substances and placed between a Crookes tube and the condenser and measuring the resulting leakage of the condenser.

I am now engaged in making a series of measurements, using this method and a condenser especially constructed for the purpose, and hope to give the results in a subsequent number.

It would seem that the method is capable of giving results much more quantitative in character than any that can be obtained by photographic methods.

WM. LISPENARD ROBB.

TRINITY COLLEGE, March 25, 1896.

AN APPARATUS FOR THE STUDY OF SOUND INTENSITIES.

The study of sound intensities presents many difficulties to the physicist as well as to the psychologist; the determination of the equality of loudness of two sounds, as well as of the law of relation between the physical cause and the sensational result is perhaps the most serious one. The facts that sounds must be estimated successively and should be of a constant intensity from beginning to end further complicate the problem. The method of the falling ball has been most frequently used; it consists in dropping a ball successively from two different heights and recording the minimum difference in height necessary to

^{*}London Electrician, February 7, 1896.